

Voice disorders are characterized by the abnormal production and/or absence of vocal quality, pitch, loudness, resonance, and/or duration, given an individual's age and/or sex.

#2

Vocal disturbances in children are surprisingly common.
A child who presents with hoarseness demands a rapid and thorough assessment
The occurrence of hoarseness ranges from 6% to 23% in school-aged children.

#3

Voice disorders in children may impede their academic performance and their socialization in school.

Moreover, the majority of adult voice disorders are thought to begin during childhood

#4

Early identification of pediatric disfluency and voice disorders is advisable because these disorders may progress to lifelong communicative impairments if left untreated.

statistics from www.asha.org



Voice disorders are characterized by the abnormal production and/or absence of vocal quality, pitch, loudness, resonance, and/or duration, given an individual's age and/or sex.

#2

Vocal disturbances in children are surprisingly common.
A child who presents with hoarseness demands a rapid and thorough assessment
The occurrence of hoarseness ranges from 6% to 23% in school-aged children.

#3

Voice disorders in children may impede their academic performance and their socialization in school.

Moreover, the majority of adult voice disorders are thought to begin during childhood

#4

Early identification of pediatric disfluency and voice disorders is advisable because these disorders may progress to lifelong communicative impairments if left untreated.

statistics from www.asha.org



Voice disorders are characterized by the abnormal production and/or absence of vocal quality, pitch, loudness, resonance, and/or duration, given an individual's age and/or sex.

#2

Vocal disturbances in children are surprisingly common.
A child who presents with hoarseness demands a rapid and thorough assessment
The occurrence of hoarseness ranges from 6% to 23% in school-aged children.

#3

Voice disorders in children may impede their academic performance and their socialization in school.

Moreover, the majority of adult voice disorders are thought to begin during childhood

#4

Early identification of pediatric disfluency and voice disorders is advisable because these disorders may progress to lifelong communicative impairments if left untreated.

statistics from www.asha.org